

Blueberry Micro Mix™

Sugar chelated micro nutrients specially designed to address common mineral deficiencies in blueberry crops. Used as a preventative when added into a fertility program.

STORAGE

Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in temperatures between 5 C (41 F) and 25 C (77 F).

Dispose of Waste Material according to local, state, and Federal environmental laws and regulations.

CAUTIONS

Because of the nature of this product care should be taken when mixing with humic acid, phosphates and or Calcium products. Always do a jar compatibility test before mixing any two products other than water in your tanks.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the user's responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water. If irritation persists, obtain medical attention.

**Ingestion: If patient is conscious, give large quantities of water. Do not induce vomiting. Call physician.*

"COMMERCIAL USE ONLY"

UAS Of America, Inc.
534 CR 529 A
Lake Panasoffkee, Florida 33538
Tel 352-793-1682 | Fax 352-793-4864
www.uasofamerica.com

GUARANTEED MINIMUM ANALYSIS

Magnesium (Mg)	3.00%
3.00% Chelated Magnesium (Mg)	
Iron (Fe)	2.000%
2.00% Chelated Iron (Fe)	
Zinc (Zn)	0.400%
0.40% Chelated Zinc (Zn)	
Manganese (Mn)	0.50%
0.50% Chelated Manganese (Mn)	
Boron as (B)	0.25%
Copper (Cu)	0.12%
Combined Sulfur	5.50%

Derived from: Magnesium Glucoheptonate, Zinc Glucoheptonate, Iron Glucoheptonate, Copper Sulfate, Manganese Glucoheptonate Complex and Sodium Borate

COMPATIBILITY

A jar compatibility test is recommended before mixing with any other product other than water in the tank to insure compatibility with foreign spray mixes and before mixing with other products other than water in the tank. Not compatible with High concentrated Calcium carbon or phosphate mixes.

APPLICATION

ORNAMENTAL AND YOUNG PLANTS: Mix 2 tsp. Blueberry Micro Mix™ with 1 gal of water. Moisten both sides of the leaves only to the point of run-off. Use misting spray bottle to apply. *Use every 7-10 days or as needed.*

AGRICULTURE: Apply 1-2 pt(s) for ROW CROPS to 2-3 qt(s) for High Bush Blueberries with sufficient water to cover per acre. Moisten both sides of the leaves only to the point of run-off.

BLUEBERRY INJECTION: Repeat as indicated visually or by leaf tissue test, or as part of a regular maintenance foliar feed program and can be applied up to - every 10 days when using the injection feeding program.

***concentrated do not over apply.**

WARRANTY

BUYER ASSUMES ALL RESPONSIBILITY FOR USE, STORAGE, HANDLING AND DISPOSAL OF THIS PRODUCT WHEN NOT IN ACCORDANCE WITH LABEL DIRECTIONS. UAS of AMERICA, INC AND ITS DISTRIBUTOR'S LIMIT OF WARRANTY EXTENDS ONLY TO REPLACEMENT OF DEFECTIVE PRODUCT.