

## COPPER X™

Copper X™ is used to correct pre existing copper deficiencies found in plants and soil. We sugar chelate our copper to readily absorb into the Plant.

## STORAGE

Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in temperatures between 5 C (41 F) and 25 C (77 F).

**Dispose of Waste Material according to local, state, and Federal environmental laws and regulations.**

## CAUTIONS

**WARNING:** Do not mix with low pH solutions or high calcium or phosphate solutions. Copper X™ will activate other materials if added to mix and cause possible burn. Do not mix with other products. Preferably spray Copper X™ alone in water mix.

Copper X™ should not be sprayed a week before or after any spray oil application.

## MIXING INSTRUCTIONS

Copper X™ is not recommended to be mixed with pesticides, Herbicides or chlorine fertilizers.

A jar compatibility test is recommended. Follow this mixing sequence:

1. Water
2. Copper X™

## DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the user's responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water. If irritation persists, obtain medical attention.

*\*Ingestion: If patient is conscious, give large quantities of water. Do not induce vomiting. Call physician.*

"COMMERCIAL USE ONLY"

UAS Of America, Inc.  
534 CR 529 A  
Lake Panasoffkee, Florida 33538  
Tel 352-793-1682 | Fax 352-793-4864  
www.uasofamerica.com

## GUARANTEED MINIMUM ANALYSIS

Copper (Cu)	5.000%
5.00% Chelated Copper (Cu)	

Derived from: Copper Glucoheptonate Chelate. Each gallon contains 0.5 lbs of elemental copper

## COMPATIBILITY

A jar compatibility test is recommended before mixing with any product other than water in the tank to insure compatibility with foreign spray mixes. Not compatible with High concentrated Calcium, carbon, phosphate or low pH mixes or spray oils. It is advisable not to mix with other materials when applying.

## APPLICATION

**FRUITBEARING:** (blueberry, cranberry, strawberry, blackberry) mix 1-2 pt(s) Copper X™ with sufficient water to cover, per acre. Application should be done prior to bloom, DO NOT apply after fruit set.

**FIELD CROPS:** (cotton, peanuts, sorghum, soybeans, and sugar beets)

**VEGETABLE CROPS:** (lettuce, melons, carrots, onions potatoes, and tomatoes) mix 1-4 pt(s) with sufficient water to cover, per acre. Apply during appropriate stage in crops cycle. DO NOT apply when fruit has formed on crop. Apply during appropriate stage in crops cycle, first application when at maturing stage.

**TREE CROPS:** Mix up to 3qts with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. USE A MINIMUM OF 100 GALLONS OF WATER.

**TURF:** Mix 2.5 oz per 1,000 sq. ft. with sufficient water to cover. USE A MINIMUM OF 15 GALLONS OF WATER.

**Apply when Copper deficiency is determined by visual or leaf analysis.**

## WARRANTY

BUYER ASSUMES ALL RESPONSIBILITY FOR USE, STORAGE, HANDLING AND DISPOSAL OF THIS PRODUCT WHEN NOT IN ACCORDANCE WITH LABEL DIRECTIONS. UAS AMERICA, INC AND ITS DISTRIBUTOR'S LIMIT OF WARRANTY EXTENDS ONLY TO REPLACEMENT OF DEFECTIVE PRODUCT.