

GUARANTEED ANALYSIS:

Magnesium (Mg) 3.00% Chelated Magnesium (Mg)	3.00%
Iron (Fe)	2.00%
2.00% Chelated Iron (Fe)	
Zinc (Zn)	0.40%
0.40% Chelate Zinc (Zn)	
Manganese (Mn)	0.50%
0.50% Chelated Manganese (Mn)	
Boron as (B)	0.25%
Copper as (Cu)	
Combined Sulfur as (S)	5.50%
Derived from: Magnesium Glucohep	tonate, Zinc
Glucoheptonate, Iron Glucoheptonate	e, Copper Sulfate,
Manganese Glucoheptonate Complex	k and Sodium Bora

WARNING

Overapplication or excessive use of boron, or use on crops other than those listed, can result in boron toxicity. This may cause foliar damage or serious crop injury.

DIRECTIONS FOR USE:

Ornamental & Young Plants: Mix 2 tsp. Blueberry Micro Mix with 1 gal of water. Moisten both sides of the leaves only to the point of run-off. Use misting spray bottle to apply. *Use every 7-10 days or as needed.*

AGRICULTURE: Apply 1-2 pt(s) for ROW CROPS to 2-3 gt(s) for High Bush Blueberries with sufficient water to cover per acre. Moisten both sides of the eaves only to the point of run-off.

Blueberry Injection

Repeat as indicated visually or by leaf tissue test, or as part of a regular maintenance foliar feed program. *Can be applied up to every 10 days when using the injection feeding program.*

*DO NOT mix with other products. Highly concentrated DO NOT over apply.

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating. "NEVER MIX CONCENTRATES DIRECTLY TOGETHER

Net Content: Size / Weight

2.5 ql/26.85 lbs 5 ql/53.7 lbs 30 ql/322.2 lbs 55 ql/ 590.7lbs 275 ql/2953.5lbs

Borate.

Blueberry Micro Mix[®]

Complex Micro-Nutrients

FOR COMMERCIAL USE ONLY

PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fundicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be spraved with this mix of chemicals to determine compatibility of products and compatibility with crop(s) spraved. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

CAUTION: KEEP OUT OF REACH OF CHILDREN



534 CR 529A Lake Panasoffkee, FL 33538 (352)-793-1682 www.uas-cropmaster.com F001142

Storage: Keep between 5 C (icro nce the ingredients are 100% foliar spray sting not rep Ξ. agnesium, Manganese ⁄/g, Mn, Zn, Fe, Mo, S soluble and immediate where മ ∠inc, li Cu and Zinc, Iron, Sume Ju and B deficiencies 3 micr 5 nutrients Copper and plant as ind Result (wnen ≤a ind Boron used are ar application as directed), evident leaf designed than 0 this ne ā s product is e to two days d for soil and/or s f of higher soil test õ

NICro

sealed tightly

wnen

not in use.

. Do

expose

d

sunlight for long

periods of time

Store in temperatures



application

Product #