

CaBN

GUARANTEED ANALYSIS:

Total Nitrogen (N)	7.00%
7.00% Ammoniacal Nitrogen	
Available Phosphate (P2O5)	0.00%
Soluble Potash (K2O)	0.00%
Calcium (Ca)	4.50%
Boron (B)	1.00%

WARNING

Overapplication or excessive use of boron, or use on crops other than those listed, can result in boron toxicity. This may cause foliar damage or serious crop injury.

PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Seller. In no case shall Seller or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. Seller makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

DIRECTIONS FOR USE:

Foliar Application Rates: Mix 1-2 quarts with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. Apply when Calcium deficiency is determined by visual or leaf analysis.

Soil Application Rates: Mix 5-11 quarts with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only Repeat every 7-10 days or as needed, until harvest. (USE MINIMUM OF 15 GALLONS OF WATER.)

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying products while tank is circulating.

"NEVER MIX CONCENTRATES DIRECTLY TOGETHER "

CAUTION: KEEP OUT OF REACH OF CHILDREN

Manufactured by:



534 CR 529A
Lake Panasoffkee, FL 33538
(352)-793-1682

www.uas-cropmaster.com

Net Content: Size / Weight

2.5 g/ 25.6 lbs 5 g/ 51.1 lbs 30 g/ 306.6 lbs 55 g/562.1 lbs 275 g/2810.5 lbs