



COPPER X™

CHELATED LIQUID COPPER MIX

FOR COMMERCIAL USE ONLY

GUARANTEED ANALYSIS:

COPPER (Cu)..... 5.00%
5.00% Chelated Copper (Cu)

Derived from: Copper Glucoheptonate Chelate.
Each gallon contains 0.5 lbs of elemental copper.
F001142

DIRECTIONS FOR USE:

DO NOT APPLY WITHIN 4 WEEKS of ANY CROP OIL APPLICATIONS

FRUIT BEARING: (blueberry, cranberry, strawberry, blackberry) mix 1-2 pt(s) CopperX with sufficient water to cover, per acre. Application should be done prior to bloom, DO NOT apply after fruit set.

FIELD CROPS: (cotton, peanuts, sorghum, soybeans, and suagrbeats) **VEGETABLE CROPS:** (lettuce, melons, carrots, onions potatoes, and tomatoes) mix 1-4 pt(s) with sufficient water to cover, per acre. Apply during appropriate stage in crops cycle. **DO NOT** apply when fruit has formed on crop. Apply during appropriate stage in crops cycle. 1st application when at maturing stage.

Tree Crops : Mix up to 3qts with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off.
 (USE MINIMUM OF 100 GALLONS OF WATER.)

Turf: Mix 2.5 ozs per 1,000 sq. ft. with sufficient water to cover. (USE MINIMUM OF 15 GALLONS OF WATER.)

Apply when Copper deficiency is determined by visual or leaf analysis.

Mixing procedure:

- 1 Fill half of spray tank with water.
- 2 Agitate water while adding COPPERX

It is UAS's recommendation that COPPERX should be applied alone in tank. If consumer desires to add COPPERX with other product(s) ie fertilizer or pesticides. It is recommended that a small jar test is mixed first and applied to the surface of several plants. **WAIT SEVERAL DAYS** to measure outcome, before proceeding with entire crop application.

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating.
 *NEVER MIX CONCENTRATES DIRECTLY TOGETHER *

Net Content: Size / Weight

2.5 gl/24.3 lbs 5 gl/48.6 lbs 30 gl/291.60 lbs 55 gl/534.60 lbs 275 gl/2,673 lbs

PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Manufactured by:



534 CR 529A
 Lake Panasoffkee, FL 33538
 (352)-793-1682
 www.uas-cropmaster.com

COPPERX is a chelated COPPER Source, designed for soil and/or foliar application, to prevent and correct pre-existing deficiencies as indicated by plant leaf, visual and/or soil testing. Since the ingredients in COPPERX are 100% water soluble and immediately available to the plant (When used as directed), this product is extremely efficient as a foliar spray in situations where a rapid response is desired. Research has shown the plant can assimilate micro nutrients better via foliar application than if the micro nutrients where applied to the soil.

Storage: Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in temperatures between 5 C (41 F) and 25 C (77 F).



Product # 50

Dispose of Waste Material according to local, state, and federal environmental laws and regulations.