



## FOR COMMERCIAL USE ONLY

#### PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eves, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

#### DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

#### CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

plant can assim esponse is quicker wnere water soluble e-existing a rapid response Mg, Mn, Fe, and chelated Magnesium, Manganese, and immediately ate micro and seems to 5 nutrients desired S available to the deficienc be more Results cies as ≤a effective are foliar plant (when Iron, and Sulfur Source, desigindicated by plant leaf, visual, often application evident used one than as đ if the directed) Ť designed for days after and/or soil this nutrie product application . soil testing. where applied to and/or foliar application, testing. Since the ingred <u>5</u>. extremely Research efficient the ingred the has shown soil. as ₫ ients മ ð prevent and correct Б П spray are S

container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in

temperatures Storage: Keep

between 5

C

(41

and

25

C (77

П



# 100%

roduct # 33

Manufactured by: of America

534 CR 529A Lake Panasoffkee, FL 33538 (352)-793-1682 www.uas-cropmaster.com F001142

### **GUARANTEED ANALYSIS:**

Magnesium (Mg) 1.00% Chelated Magnesium (Mg)	1.00%
Iron (Fe)	3.00%
3.00% Chelated Iron (Fe)	
Manganese (Mn)	4.00%
4.00% Chelated Manganese (Mn)	
Sulfur (S)	5.50%
Derived from: Magnesium Glucoheptonate, Iron	
Glucoheptonate, Manganese Glucoheptonate.	

#### DIRECTIONS FOR USE:

Ornamental & Young Plants: Mix 2 tsp. F-1 Mix with 1 gal of water. Moisten both sides of the leaves only to the point of run-off. Use misting spray bottle to apply. \*Use every 7-10 days or as needed.\*

Gardens, Young Trees & Hardy Plants: Mix 1 tbsp. F-1 Mix to each gal, of water, Moisten both sides of the leaves only to the point of run-off. Use misting spray bottle or pump up spraver to apply. \*Use every 10-14 days or as needed. \*

Lawns: Apply 1qt per 10,000 sq. ft. with hose end sprayer. Set dial on 2 oz/gal. Apply before or after each fertilization. \*Use only 1-2 times per season. \*

Hardy Trees: Mix 1/2 to 1 oz F-1 Mix to each 1 gal of water. Moisten both sides of the eaves only to the point of run-off. Use misting spray bottle or pump up sprayer to apply.

AGRICULTURE: Apply 1-2 pt. for ROW CROPS to 2-3 pt. for FRUIT TREES, F-1 Mix with sufficient water to cover per acre. Moisten both sides of the leaves only to the point of run-off.

Greens & Tees: Use 6 oz per 1,000 sq. ft. Once a month. Fairways: Use 4 oz. per 1,000 sg. ft. Once a month.

Repeat as indicated visually or by leaf tissue test, or as part of a regular maintenance foliar feed program. \*

\*\*\*Highly concentrated DO NOT over apply. \*\*\* To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating "NEVER MIX CONCENTRATES DIRECTLY TOGETHER "

CAUTION: KEEP OUT OF REACH OF CHILDREN

55 gl/ 605 lbs

275 al/ 3.025 lbs

Net Content: Size / Weight

2.5 ql/ 27.5 lbs 5 ql/ 55 lbs 30 ql/ 330 lbs

Mix is Do I need a F-1 Mix?