

GUARANTEED ANALYSIS:

Magnesium (Mg)	1.70%
1.70% Chelated Magnesium (Mg)	
Zinc (Zn)	0.75%
0.75% Chelated Zinc (Zn)	
Iron (Fe)	3.50%
3.50% Chelated Iron (Fe)	
Manganese (Mn)	0.75%
0.75% Chelated Manganese (Mn)	
Boron (B)	0.20%
Molybdenum (Mo)	0.003%
Combined Sulfur (S)	4.00%
Derived from: Magnesium Glucohepte	onate. Zinc Glu

ucoheptonate. Iron Glucoheptonate, Sodium Molybdate, Manganese Glucoheptonate Complex and Sodium Borate.

Non-Plant Food Ingredient:

Alpha Keto Acid.....0.02% Purpose: To enhance nutrient uptake.

DIRECTIONS FOR USE:

AGRICULTURE: Apply 1-2 pt(s) for ROW CROPS to 2-3 gt(s) for FRUIT TREES, MICRO BOOST with sufficient water to cover per acre. Moisten both sides of the leaves only to the point of run-off.

*Repeat as indicated visually or by leaf tissue test, or as part of a regular maintenance foliar feed program. *

*DO NOT mix with other products, Highly concentrated DO NOT over apply.

Ornamental & Young Plants: Mix 2 tsp. MICRO BOOST with 1 gal of water. Moisten both sides of the leaves only to the point of run-off. Use misting spray bottle to apply. *Use every 7-10 days or as needed.*

LAWNS/COMMERCIAL TURF: Apply 3 oz per 1000 sg. ft. with hose end. Set dial on 2oz/gal. Apply before or after each fertilization. *Use min 2 times per season. *

*** Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating. "NEVER MIX CONCENTRATES DIRECTLY TOGETHER

Net Content: Size / Weight

2.5 gl/ 26.85lbs 5 gl/53.7 lbs 30 gl/322.2 lbs 55 gl/590.7 lbs 275 gl/ 2953.5lbs

MICRO BOOST ®

Chelated Micro-Nutrients & Alpha Keto Acid

FOR COMMERCIAL USE ONLY PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eves, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.


CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential. special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

WARNING

Overapplication or excessive use of boron, or use on crops other than those listed, can result in boron toxicity. This may cause foliar damage or serious crop injury.

This fertilizer contains molybdenum (Mo). The application of fertilizer materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant



534 CR 529A Lake Panasoffkee, FL 33538

(352)-793-1682 www.uas-cropmaster.com

F001142

Storage: Keep between 5 C (micro product atter appli) nutrients extremely (41 F) and where applied to the aner sealed tightly v ⁻) and 25 C (77 F). efficient Rese as മ foliar spray not the Ξ t in use. e plant can S. s quicker s where a rapid n assimilate mici and licro diately available : d response is des icro nutrients bett sunlight for long 5 more s desired. better via periods of time ₫ (when ults are r applic Store in temperatures ofter ten d as directed) n evident one n than if the nt one to the

Why Do I need Micro Boost? MICRO BOOST is a chelated Magnesium, application, to prevent and correct pre-exis Since the ingredients in Micro-Boost are 10

pre-existing

, Mangar sting Mg,

100%

ē

Immed

and

nd B c liately r, Molybdenum anu nd B deficiencies a -:Iable to f

as i

Sulfur, S

and

Boron na

ion Source, dicated by p

designed for

by plant

leaf

visual

-e, N and e, Mo

soil and/or folia al and/or soil te ¥ testing



Dispose of Waste Material according to local, state, and federal environmental laws and regulations