



Micro Hume[®]

Trace elements with organic Substance activators

FOR COMMERCIAL USE ONLY

PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

WARNING

Overapplication or excessive use of boron, or use on crops other than those listed, can result in boron toxicity. This may cause foliar damage or serious crop injury.

This fertilizer contains molybdenum (Mo). The application of fertilizer materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant

Manufactured by:



534 CR 529A
Lake Panasoffkee, FL 33538
(352)-793-1682

www.uas-cropmaster.com
F001142

Why Do I need MICRO HUME?
Research has shown Humic - Fulvic acids to be one of the best natural chelating agents available. This unique combination of the Humic / Fulvic acid with microelements allows for vigorous stimulating effects. The unique combination of compounds found in Hume - Fulvic acids combined in Micro-Hume with it's Micro-Elements are involved in the DNA activation in the nucleus of plant cells. Also affecting the RNA within a cell, this combination acts as a catalyst helping to speed up the rate of chemical reactions within plant cells. With added carbohydrates this combination allows for complete absorption with additional growth and cell stimulation. Micro-Hume will help activate enzymes and intensify the metabolism of proteins having a positive growth regulating effect on plants. This will also help to regulate rapidly changing hormone levels of plants going through stress levels IE Drought, Wet, Cold, Heat, Flowering and transplanting.

Storage: Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time.



Product # 86

GUARANTEED ANALYSIS:

Magnesium (Mg).....	1.50%
1.50% Chelated Magnesium (Mg)	
Zinc (Zn).....	0.50%
0.50% Chelated Zinc (Zn)	
Iron (Fe).....	3.00%
3.00% Chelated Iron (Fe)	
Manganese (Mn).....	0.80%
0.80% Chelated Manganese (Mn)	
Copper (Cu).....	0.01%
0.01% Chelated Copper (Cu)	
Boron (B).....	0.50%
Molybdenum (Mo).....	0.01%

Derived from: Magnesium Glucoheptonate, Zinc Glucoheptonate, Iron Glucoheptonate, Copper Glucoheptonate, Manganese Glucoheptonate Complex, Sodium Molybdate, and Sodium Borate.

Non-Plant Food Ingredient:

0.70% Fulvic Acid	Alph Keto Acid	0.02%
Derived from: Leonardite Shale.	Purpose: To enhance nutrient uptake	

DIRECTIONS FOR USE:

Vegetable/Row Crops: Apply 2 qts. per acre with band sprayer. Apply only to point of run off. Cover entire plant.

Blueberry/Blackberry: Apply on a foliar application rate of 1-2 qt(s). per acre. Apply every 30 days from March-to October. Or apply as needed.

Citrus: Apply Micro Hume during post-bloom at a rate of 3 qts. per acre. Repeat if necessary during summer oil spray.

Tropical Fruits, Persimmons and other Deciduous Fruits, Pecans and other Nuts: Apply Micro Hume in the late bloom to pedal drop stage at 3 qts. per acre and if necessary again in the late spring or during summer spray at 3 qts. per acre.

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating.
"NEVER MIX CONCENTRATES DIRECTLY TOGETHER"

Net Content: Size / Weight

CAUTION: KEEP OUT OF REACH OF CHILDREN

2.5 g/26.88 lbs 5 g/53.75 lbs 30 g/322.5 lbs 55 g/591.25 lbs 275 g/2,956.25 lbs

Dispose of Waste Material according to local, state, and federal environmental laws and regulations.